

Fermi-Bose Mixtures Near Broad Interspecies Feshbach Resonances

Jun Liang Song, Mohammad S. Mashayekhi and Fei Zhou

Department of Physics and Astronomy, The University of British Columbia, Vancouver, B. C., Canada V6T1Z1

(Dated: June 5, 2010)

In this Letter we have studied dressed bound states in Fermi-Bose mixtures near broad interspecies resonance, and implications on many-body correlations. We present the evidence for a first order phase transition between a mixture of Fermi gas and condensate, and a fully paired mixture where extended fermionic molecules occupy a single pairing channel instead of forming a molecular Fermi surface. We have further investigated the effect of Fermi surface dynamics, pair fluctuations and discussed the validity of our results.

Since the observations of molecules of Fermi atoms near Feshbach resonances, fascinating pairing correlations in cold Fermi gases have been successfully investigated both experimentally [1–5] and theoretically [6–8]. Near broad resonances where the atom-molecule coupling is very strong, pair correlations can also be closely related to the ones in the BEC-BCS crossover theory pioneered a while ago [9–11]. Meanwhile, interspecies Feshbach resonances in Fermi-Bose mixtures of ${}^6\text{Li}$ - ${}^{23}\text{Na}$, ${}^{40}\text{K}$ - ${}^{87}\text{Rb}$ and ${}^6\text{Li}$ - ${}^{87}\text{Rb}$ have been experimentally observed [12–16]. Recently, weakly bound ${}^{40}\text{K}$ - ${}^{87}\text{Rb}$ pairs prepared near Feshbach resonance were further successfully converted into cold molecules at JILA [17], which can potentially lead to exciting opportunities of studying new quantum states of matter [18]. Previous theoretical studies on Fermi-Bose mixtures on the other hand have been mainly focused on narrow resonances or when the atom-molecule coupling is very weak [19–21]; phase boundaries in this limit depend on atom-molecule coupling strengths.

Experimentally, creating and probing correlations in Fermi-Bose mixtures near narrow resonances are more challenging than near broad resonances. For a Feshbach resonance with a width ΔB and background scattering length a_{BG} , an effective resonance energy width can be introduced as $\Gamma_{res} = 2m_R a_{BG}^2 (\Delta\mu \Delta B)^2 / \hbar^2$ [10, 22]; here $\Delta\mu$ is the difference in magnet moments between the scattering and molecule channel and m_R is the reduced mass of a pair of Bose and Fermi atoms. For interspecies resonances so far confirmed [12–16, 23], Γ_{res} can be a few orders of magnitude bigger than the quantum degeneracy energy at a density of 10^{14}cm^{-3} [12]. So some of well studied resonances are quite broad and many-body correlations near these interspecies resonances remain to be thoroughly understood. One of very fundamental questions we hope to answer in this Letter is *what is the nature of quantum matter near broad interspecies resonances*. Especially, how do dressed two-body bound states evolve when approaching resonances? and accordingly what kind of many-body correlations are developed in a quantum Fermi-Bose mixture? And physics near broad resonances can distinctly differ from that near narrow resonances; some basic concepts introduced for narrow resonances such as molecular Fermi surfaces might not be directly applicable here. Motivated by these con-

siderations, we carry out a study on Fermi-Bose mixtures near broad interspecies Feshbach resonances which serves as a potential reference for more sophisticated analyses. Apart from the phase diagrams in terms of interspecies scattering lengths a_{bf} and Bose(m_B)-Fermi(m_F) mass ratio m_B/m_F , we focus on bound state properties of near-resonance Fermi-Bose mixtures which can be potentially probed in experiments. Our results are useful for the understanding of correlations in ${}^6\text{Li}$ - ${}^{23}\text{Na}$, ${}^6\text{Li}$ - ${}^{87}\text{Rb}$, and ${}^{40}\text{K}$ - ${}^{87}\text{Rb}$ mixtures. For simplicity, we employ a simplest one-channel Hamiltonian which captures most important aspects near broad resonances,

$$H = \sum_{\mathbf{k}} \epsilon_{\mathbf{k}}^F f_{\mathbf{k}}^\dagger f_{\mathbf{k}} + \sum_{\mathbf{k}} \epsilon_{\mathbf{k}}^B b_{\mathbf{k}}^\dagger b_{\mathbf{k}} + \frac{V_{bf}}{\Omega} \sum_{\mathbf{k}, \mathbf{k}', \mathbf{Q}} f_{\frac{m_B}{m_F} \mathbf{Q} + \mathbf{k}}^\dagger b_{\frac{m_B}{m_F} \mathbf{Q} - \mathbf{k}}^\dagger f_{\frac{m_B}{m_F} \mathbf{Q} + \mathbf{k}'} b_{\frac{m_B}{m_F} \mathbf{Q} - \mathbf{k}'} \quad (1)$$

where $f_{\mathbf{k}}^\dagger, b_{\mathbf{k}}^\dagger$ ($f_{\mathbf{k}}, b_{\mathbf{k}}$) are creation (annihilation) operators for Fermi and Bose atoms respectively, and $\epsilon_{\mathbf{k}}^{F(B)} = \frac{\hbar^2 |\mathbf{k}|^2}{2m_{F(B)}}$ are kinetic energies for fermions (bosons) and Ω is the volume. V_{bf} is the strength of interaction that is related to interspecies scattering lengths a_{bf} via

$$\frac{1}{V_{bf}} = \frac{m_R}{2\pi a_{bf} \hbar^2} - \frac{1}{\Omega} \sum_{\mathbf{k}} \frac{1}{\epsilon_{\mathbf{k}}^R}; \quad (2)$$

here $m_R = m_B m_F / (m_B + m_F)$, $\epsilon_{\mathbf{k}}^R = \hbar^2 \mathbf{k}^2 / 2m_R$. We assume that the background boson-boson interactions are repulsive so that the mixture is stable; to illustrate the idea, here we only include interspecies scattering.

Bound states and Pauli blocking effects We first consider the binding energy of a pair of Fermi and Bose atoms with opposite momenta ($\mathbf{k}, -\mathbf{k}$), in the presence of a condensate (BEC) and a Fermi surface of Fermi atoms which blocks all states below its Fermi momentum $\hbar k_F$. Pauli blocking effects of a Fermi sea indeed lead to *dressed* bound states at arbitrary small negative scattering lengths but with an anomalous dispersion, or a negative effective mass (see also discussions before Eq.4). Furthermore, the energy W_B it takes to create a bound state from a non-interacting ground state




FIG. 1. (color online) The energy dispersion of bound states for mass ratio $m_B/m_F = 2.175$ or ^{40}K - ^{87}Rb mixtures. From the top to bottom are $W_B(\mathbf{Q})$ in units of ϵ_F^R for $1/k_F a_{bf} = -0.2, -0.1145, -0.05, 0.0145$. The shaded region represents pair excitation continuum.

can be either positive or negative depending on scattering lengths a_{bf} , a unique feature of Fermi-Bose systems. This is because to form a pair of atoms with opposite momenta $(\mathbf{k}, -\mathbf{k})$ near the Fermi surface $|\mathbf{k}| = k_F$, a Bose atom has to be promoted to right above the Fermi surface which results in an energy penalty $\epsilon_F^B = \hbar^2 k_F^2 / 2m_B$. For a bound state with an arbitrary total momentum $\mathbf{Q}$ or a kinetic energy $\epsilon_Q^C = \hbar^2 \mathbf{Q}^2 / 2(m_F + m_B)$, the energy cost is $W_B(\mathbf{Q}) = \epsilon_F^B + \epsilon_Q^C + \omega_B$. The $\mathbf{Q}$ -dependent binding energy $\omega_B (< 0)$ can be obtained by solving the following two-body equation,

$$\frac{-m_R \Omega}{2\pi a_{bf} \hbar^2} = \left(\sum_{|\frac{m_B}{m_F} \mathbf{Q} + \mathbf{k}| > k_F} \frac{1}{\epsilon_{\mathbf{k}}^R - \epsilon_F^R - \omega_B} - \sum_{\mathbf{k}} \frac{1}{\epsilon_{\mathbf{k}}^R} \right). \quad (3)$$

In the limit of small $k_F a_{bf} (< 0)$ and when $\mathbf{Q} = 0$, Eq.(3) leads to $\omega_B = -4\epsilon_F^R \exp(\frac{\pi}{k_F a_{bf}})$, $\epsilon_F^R = \hbar^2 k_F^2 / 2m_R$. The dispersion of bound states that can be probed using photoassociative spectroscopy is shown in Fig.1 [24]. For small negative scattering lengths a_{bf} , bound states are fully gapped with positive energies $W_B(\mathbf{Q})$, and the ground state is a mixture of Fermi gas and BEC. However, $W_B(0)$, the energy gap of bound states vanishes at a *critical* scattering length $a^{(1)}$. In Fig.2, we present results of $a^{(1)}$ versus m_B/m_F . For heavy Bose atoms, $k_F a^{(1)}$ approaches a small value of $\pi / (\ln[m_F / 4m_B] + 2)$.

To ensure the stability of $\mathbf{Q} = 0$ molecules near the transition line $a^{(1)}$, we further examine M_{eff} , the effective mass near $\mathbf{Q} = 0$. At scattering lengths $a^{(1)}$ or when $W_B(0) = 0$, we find

$$\frac{1}{M_{eff}} = \frac{1}{m_T} \left[1 - \frac{4m_F}{3m_B} g\left(\frac{m_R}{m_F}\right) \right]; \quad (4)$$




FIG. 2. (color online) Scattering length $a^{(1)}$ at which the energy cost W_B to create a molecule with $\hbar \mathbf{Q} = 0$ from a Fermi gas-BEC mixture becomes zero, as a function of Bose-Fermi mass ratio m_B/m_F . Inset is W_B , the energy gap of molecules in units of ϵ_F^R as a function of $k_F a_{bf}$ (and $1/a_{bf} < 1/a^{(1)}$) for different mass ratios or mixtures.

and the dimensionless function $g(x^2) = x / [(1 - x^2)^2 (\ln \frac{1+x}{1-x} + \frac{2x}{1-x^2})]$. As far as $m_B/m_F > 0.7$ and the energy penalty ϵ_F^B is not too heavy, M_{eff} is positive, although it can be are much bigger than the bare mass $m_T (= m_F + m_B)$ as a result of dressing in the Fermi sea. Below we focus on the limit of positive M_{eff} that is most relevant to the experimental mass ratio m_B/m_F (between 2.175 and 14.5) [24]. Note that the binding energy ω_B is independent of the Bose atom density when the Fermi sea is treated as a static background.

The above analysis at first sight seems to suggest that when W_B becomes negative, a small fraction of Fermi and Bose atoms start forming molecules or a dilute molecular Fermi gas signifying a phase transition at $a^{(1)}$. Such a picture was in fact previously proposed for mixtures near *narrow* resonances [19, 20]. However since the extent of molecules d_m is typically comparable to or much longer than the Fermi wave length $2\pi/k_F$ near broad resonances, pairs may be accommodated, even before the two-body gap W_B vanishes, in other more exotic forms without forming a molecular Fermi surface. Below we provide evidence for such a possibility. Note that a finite two-body gap W_B suggests a local stability of the Fermi gas-BEC mixture against emergence of a Fermi gas of molecules, i.e. when $1/a < 1/a^{(1)}$, a molecular Fermi gas can *not* be a ground state. It is in this limit that we illustrate, based on an energetic analysis, that a *third* state or a fully paired state actually further lowers the energy.

Energy landscape of pairing states Suggested by above discussions, we consider energetics of pairing states of Fermi-Bose atoms and from now on focus on *homogeneous* mixtures with equal populations of fermions and bosons, i.e. $N_F = N_B$. Although in principle pairing with a finite total momentum $\hbar \mathbf{Q}$ can occur in ground

states, detailed calculations show that for a range of mass ratios ($m_B/m_F > 0.2$) that is relevant to Fermi-Bose mixtures so far studied in experiments, pairing in $\hbar\mathbf{Q} = 0$ channel is always dominating and favored, qualitatively consistent with our analyses on the two-body bound states. Below we only show results of $\mathbf{Q} = 0$ pairing and adopt the simplest pairing wavefunction

$$|g.s.\rangle = \exp(c_0 b_0^\dagger) \prod_{\mathbf{k} \neq 0} (u_{\mathbf{k}} + v_{\mathbf{k}} f_{\mathbf{k}}^\dagger b_{-\mathbf{k}}^\dagger + \eta_{\mathbf{k}} f_{\mathbf{k}}^\dagger) |vac\rangle \quad (5)$$

where $u_{\mathbf{k}}$, $v_{\mathbf{k}}$ and $\eta_{\mathbf{k}}$ are three families of variational parameters. We obtain the energy of the variational states and then minimize it with respect to $u_{\mathbf{k}}$, $v_{\mathbf{k}}$ and $\eta_{\mathbf{k}}$ that are subject to the normalization condition $|u_{\mathbf{k}}|^2 + |v_{\mathbf{k}}|^2 + |\eta_{\mathbf{k}}|^2 = 1$. Equilibrium conditions can then be obtained and there are two solutions for any given $\mathbf{k}$: i) a unpaired state with $\eta_{\mathbf{k}} = 1$ and $u_{\mathbf{k}} = v_{\mathbf{k}} = 0$; ii) a paired state with $\eta_{\mathbf{k}} = 0$ and $v_{\mathbf{k}}^2 = \frac{1}{2} \left(1 - \frac{\xi_{\mathbf{k}}^R}{E_{\mathbf{k}}}\right)$, $u_{\mathbf{k}}^2 = \frac{1}{2} \left(1 + \frac{\xi_{\mathbf{k}}^R}{E_{\mathbf{k}}}\right)$, $\Delta = -\frac{V_{bf}}{\Omega} \sum_{\mathbf{k}} (1 - \eta_{\mathbf{k}}^2) u_{\mathbf{k}} v_{\mathbf{k}}$, where $E_{\mathbf{k}} = \sqrt{(\xi_{\mathbf{k}}^R)^2 + 4\Delta^2}$, $\xi_{\mathbf{k}}^R = \epsilon_{\mathbf{k}}^R - \mu$ and μ is the pair chemical potential. Pairing gap Δ , μ as well as the condensed population $|c_0|^2$ are determined self-consistently, $N_F = \sum_{\mathbf{k}} \left(v_{\mathbf{k}}^2 (1 - \eta_{\mathbf{k}}^2) + \eta_{\mathbf{k}}^2\right)$, $N_B = |c_0|^2 + \sum_{\mathbf{k}} v_{\mathbf{k}}^2 (1 - \eta_{\mathbf{k}}^2)$ and




FIG. 3. (Color online). (a) a_{cr} at which a first order phase transition occurs, as a function of mass ratio m_B/m_F . The dashed line is $a^{(1)}$ along which a Fermi gas-BEC mixture becomes locally unstable. Near a_{cr} , there is a tiny window for phase separation (shaded area in the inset). Line 2 (1) is the shifted a_{cr} ($a^{(1)}$) due to quantum fluctuations (see discussions around Eq.7). (b) E_p , energy per pair of Fermi-Bose atoms in units of ϵ_F^R as a function of X , the size of Fermi surface of unpaired Fermi atoms for $m_B/m_F = 2.175$ or ^{40}K - ^{87}Rb mixture. From the top to bottom are E_p for $1/k_F a_{bf} = -5$, -0.1145 (for a_{cr}), -0.05 , 0.0145 (for $a^{(1)}$) and 0.2 . $X = 1$ corresponds to a unpaired state and $X = 0$ is for a fully paired mixture.

$$\frac{-m_R \Omega}{2\pi a_{bf} \hbar^2} = \sum_{\mathbf{k}} \frac{1 - \eta_{\mathbf{k}}^2}{\sqrt{(\epsilon_{\mathbf{k}}^R - \mu)^2 + 4\Delta^2}} - \sum_{\mathbf{k}} \frac{1}{\epsilon_{\mathbf{k}}^R}. \quad (6)$$

To minimize the energy, we further choose $\eta_{\mathbf{k}}$ to be a step function, $\eta_{\mathbf{k}} = 1$ if $|\mathbf{k}| \leq X k_F$, and zero otherwise; the dimensionless variational parameter $X \in [0, 1]$ specifies the size of residue Fermi surface of unpaired Fermi atoms. When $N_F = N_B$, one can also verify that X^3 is equal to $|c_0|^2/N_B$, i.e. the condensation fraction.

In Fig.3,4, we present the main results of variational calculations. In Fig.3, the energy per pair of atoms is shown as a function of X , the size of Fermi surface of unpaired fermions. At small and negative scattering lengths, a Fermi gas-BEC mixture is a ground state and $X_0 = 1$, $\mu = \epsilon_F^F + W_B$ ($\epsilon_F^F = \hbar^2 k_F^2/2m_F$) and $\Delta = 0$. A fully paired state with $X_0 = 0$ becomes degenerate with the Fermi gas-BEC mixture ($X_0 = 1$) at a critical scattering length a_{cr} , beyond which the paired state becomes a ground state. However, a Fermi gas-BEC mixture remains to be locally stable until scattering lengths reach the value of $a^{(1)}$ which is fully consistent with the above study of bound states. Our variational calculations suggest a first order phase transition between a Fermi gas-BEC mixture and a fully paired mixture. This later state of extended molecules is conceptually different from a Fermi gas of molecules; instead, all molecules, though fermionic in nature, occupy the same pairing channel with zero total momentum. A direct comparison of energies indicates that a fully paired mixture has lower energies than a Fermi gas of molecules provided $1/a_{bf} < 1/a^{(2)}$. For ^{40}K - ^{87}Rb mixtures, $1/k_F a^{(2)}$ is about $0.25 - 0.3$ and $1/a^{(2)} > 1/a^{(1)}$; further towards the molecular side, a paired mixture is expected to evolve into a Fermi gas of molecules.

In Fig.4, we further present the results on the pair breaking energy Δ and pair chemical potential μ . The pair breaking energy can be probed when applying rf pulses to transfer Fermi atoms to a different hyperfine spin state[5, 25] that weakly interacts with the Fermi-Bose mixture. The frequency shift in the rf spectroscopy should be $\hbar\Delta\omega(\mathbf{k}) = \frac{1}{2}(\xi_{\mathbf{k}}^R + \sqrt{|\xi_{\mathbf{k}}^R|^2 + 4\Delta^2})$. In the fully paired phase, Bose atoms are completely depleted and the Bose atom distribution $n_B(\mathbf{k})$ follows closely the Fermi atom distribution $n_F(\mathbf{k})$.

Fermi surface dynamics and pair fluctuations

At small negative scattering lengths when quantum fluctuations are weak, our mean-field analyses on two-body bound states as well as the fully paired states are asymptotically adequate. And $a^{(1)}, a_{cr}$ estimated in the limit of large mass ratio ($k_F a^{(1)} \sim \pi/(\ln[m_F/4m_B] + 2)$ in Fig.3) are quantitatively valid. However when $k_F a$ is of order of one, the fluctuations becomes substantial and the bound states can be further dressed in fluctuating particle-hole pairs; the energetic analyses are subject to




FIG. 4. (color online). (a) Pair-breaking energy Δ and (b) pair chemical potential μ in units of ϵ_F^R as a function of scattering length $k_F a_{bf}$ (near a_{cr}) for mass ratio $m_B/m_F = 2.175$ or $^{40}\text{K}-^{87}\text{Rb}$. Dashed lines are for the values of meta-stable states. The tiny window for phase separation near a_{cr} is too small to show here (see inset of Fig.3a).

corrections. To clarify this, we estimate the dominating effects of Fermi surface dynamics, i.e. the Gorkov corrections (*GMB*) in the two-body scattering vertex due to fluctuating particle-hole pairs[22, 26, 27], and the self-energy (*SE*) effect which mainly represents the effect of atomic Fermi surface smearing and the mass renormalization due to scattering by the condensate or Fermi sea. We then examine the pole structure of the propagator for a pair of Fermi-Bose atoms taking into account the vertex corrections and self-energy effects and obtain the binding energy ω_B . The relative shift caused by the *GMB* effect (the leading order term R_1) and the *SE* effect (the higher order term R_2) is given as

$$\frac{\delta \ln |\omega_B|}{\ln |\omega_B|} = R_1(k_F a_{bf}) + R_2(k_F a_{bf})^2 \ln |k_F a_{bf}| \dots \quad (7)$$

where $R_{1,2}$ both are dimensionless quantities depending on the mass ratio m_B/m_F and can be obtained numerically[28]. For large mass ratios (m_B/m_F), one finds that $R_1 \sim -\ln(m_B/m_F)/\pi$ and $R_2 \sim -2/(3\pi^2)$; and ω_B is reduced by a factor of m_F/m_B solely due to the *GMB* effect. The net reduction in the binding energy ω_B leads to a upward shift of $1/a^{(1)}$ (line 1) in Fig.3. In addition, we have estimated that, for the paired state, the amplitude of pair fluctuations $A_1 \sim (\Delta/\epsilon_F^R)^4 \sqrt{\kappa/m_R k_F}$ (κ is the compressibility of the paired mixture), and the corresponding zero point energy (in units of ϵ_F^R) per particle $A_2 \sim (\Delta/\epsilon_F^R)^4 \sqrt{m_R k_F/\kappa}$. We incorporate these quantum fluctuations (analogous to *NSR* effects[11]) which further favor pairing, and the *GMB* correction into our analysis of energetics of the paired state. For $^{40}\text{K}-^{87}\text{Rb}$ mixtures, we find that $A_1 \approx 0.01$, $A_2 \approx 0.003$ and $R_1 \approx -0.4$ and $R_2 \approx -0.1$; the *GMB* and *SE* effects appear to be more dominating. The modified critical lines obtained by extrapolating the above analyses to near resonance are shown in Fig.3 and are qualitatively consistent with the mean field ones. Data suggest that quantum fluctuations tend to enlarge the window between $1/a_{cr}$

and $1/a^{(1)}$ and further stabilize the first order phase transition. Quantum Monte Carlo simulations similar to those in Ref. [29] remain to be carried out.

In conclusion, we have examined dressed bound states and provided evidence of a new quantum state of extended molecules near broad interspecies resonances. As far as three- and higher-body correlations are insignificant, our results can be applied to understand Fermi-Bose mixtures near broad resonances. We thank Gordon Baym, Immanuel Bloch, Kirk Madison, Subir Sachdev, Joseph Thywissen, Jun Ye and Zhenhua Yu for stimulating discussions. This work is supported by NSERC (Canada) and Canadian Institute for Advanced Research. **Note added:** Upon the submission of this work, we learned that finite-temperature mixtures near broad resonance were also studied in Ref.[30].

-
- [1] C. A. Regal *et al.*, Nature **424**, 47-50 (2003); C. A. Regal *et al.*, Phys. Rev. Lett. **92**, 040403 (2004).
 - [2] K. Strecker *et al.*, Phys. Rev. Lett. **91**, 080406 (2003).
 - [3] S. Jochim *et al.*, Phys. Rev. Lett. **91**, 240402 (2003).
 - [4] M. Zwierlein *et al.*, Phys. Rev. Lett. **91**, 250401(2003).
 - [5] C. Chin *et al.*, Science **305**, 1128-1130 (2004).
 - [6] M. Holland *et al.*, Phys. Rev. Lett. **87**, 120406 (2001); E. Timmermans *et al.*, Phys. Lett. A **285**, 228 (2001); Y. Ohashi and A. Griffin, Phys. Rev. Lett. **89**, 130402 (2002).
 - [7] T. L. Ho, Phys. Rev. Lett. **92**, 090402 (2004).
 - [8] A. V. Andreev *et al.*, Phys. Rev. Lett. **93**, 130402 (2004).
 - [9] D. M. Eagles, Phys. Rev. **186**, 456(1969).
 - [10] A. J. Leggett, J. Phys.(Paris) Colloq. **41**, 7(1980); for discussions on resonance widths, see A. J. Leggett, *Quantum Liquids* (Oxford University Press, 2006).
 - [11] P. Nozieres and S. Schmitt-Rink, J. of Low Temp. Phys. **59**, 195 (1985).
 - [12] C. A. Stan *et al.*, Phys. Rev. Lett. **93**, 143001 (2004).
 - [13] S. Inouye *et al.*, Phys. Rev. Lett. **93**, 183201 (2004).
 - [14] F. Ferlaino *et al.*, Phys. Rev. A **73**, 040702 (2006).
 - [15] S. Ospelkaus *et al.*, Phys Rev Lett. **97**, 120403 (2006).
 - [16] B. Deh *et al.*, Phys. Rev. A **77**, 010701(R) (2008).
 - [17] K. K. Ni *et al.*, Science **322**, 231 (2008).
 - [18] H. P. Büchler *et al.*, Phys. Rev. Lett. **98**, 060404 (2007).
 - [19] S. Powell *et al.*, Phys. Rev. B **72**, 024534(2005).
 - [20] H. Yabu *et al.*, Physica B **329-333**, 25 (2003).
 - [21] D. C. E. Bortolotti *et al.*, J. Phys. B **39**, 189 (2006).
 - [22] C.J. Pethick and H. Smith, *Bose-Einstein condensation in dilute gases* (Cambridge University Press, 2008).
 - [23] A. Simoni *et al.*, Phys. Rev. Lett. **90**, 163202 (2003); M. Gacesa *et al.*, Phys. Rev. A **78**, 010701(R)(2008); Z. Li *et al.*, Phys. Rev. A **78**, 022710 (2008); C. Marzok *et al.*, Phys. Rev. A **79**, 012717 (2009).
 - [24] Far away from resonances or for much smaller Bose-Fermi mass ratios, the dispersion can be anomalous with a minimum at $|\mathbf{Q}| \neq 0$.
 - [25] G. Baym *et al.*, Phys. Rev. Lett. **99**, 190407 (2007).
 - [26] L. P. Gorkov and T. M. Melik-Barkhudarov, Sov. Phys. JETP **13**, 1018 (1961).
 - [27] H. Heiselberg *et al.*, Phys. Rev. Lett. **85**, 2418 (2000).

- [28] The log-divergency in Eq.7 can be removed if the compressibility of BEC is taken into account.
- [29] S. Pilati and S. Giorgini , Phys. Rev. Lett.**100**, 030401 (2008).
- [30] E. Fratini and P. Pieri, Phys. Rev. A **81**,051605(2010).